

資料論文  
Data paper

Point count sampling data  
from the Taiwan Breeding Bird Survey

臺灣繁殖鳥類大調查的定點計數數量資料

Jerome Chie-Jen Ko<sup>1,2</sup>, Meng-Wen Fan<sup>1,\*</sup>, Ruey-Shing Lin<sup>1</sup>, Pei-Fen Lee<sup>2</sup>,  
Shih-Peng Tsai<sup>3</sup>

柯智仁<sup>1,2</sup> 范孟雯<sup>1,\*</sup> 林瑞興<sup>1</sup> 李培芬<sup>2</sup> 蔡世鵬<sup>3</sup>

<sup>1</sup> Endemic Species Research Institute, No. 1, Ming-shen East Rd., Chichi Township, Nantou, 55244, Taiwan

<sup>2</sup> Institute of Ecology and Evolutionary Biology, National Taiwan University, No. 1, Sec. 4, Roosevelt Rd., Taipei, 10617, Taiwan

<sup>3</sup> Chinese Wild Bird Federation, 1F, No. 3, Lane 36, Jing-Long St., Taipei, 11683, Taiwan

<sup>1</sup> 行政院農業委員會特有生物研究保育中心 南投縣集集镇民生東路1號

<sup>2</sup> 國立臺灣大學生態學與演化生物學研究所 臺北市羅斯福路四段1號


<sup>3</sup> 中華民國野鳥學會 台北市景隆街36巷3號

\*Corresponding author: mwluna0927@gmail.com

\*通訊作者：mwluna0927@gmail.com

## Abstract

The Taiwan Breeding Bird Survey (BBS Taiwan) is a nationwide monitoring project with the aim of establishing a breeding bird population index for conservation. The project was initiated in 2009 as a citizen science project by a cross-organizational working group. The BBS Taiwan has obtained 358,183 records of bird sampling data from 454 sampling sites during the breeding season from 2009 to 2015. This paper provides the descriptions of sampling design, survey methods, and coverage of species of the dataset.



## 摘要

臺灣繁殖鳥類大調查（簡稱BBS Taiwan）是以建立繁殖鳥類族群指標趨勢為目標的全國性監測計畫。自2009年起，透過結合政府組織、學術機關，以及民間團體的夥伴關係持續推動。累計至2015年，BBS Taiwan已累計454個樣區，358,183筆繁殖鳥類取樣事件紀錄。本篇資料論文說明計畫的取樣設計、調查方法，以及涵蓋的鳥種範圍。

**Keywords:** Sampling event, Aves, distance sampling, population index, citizen science

**關鍵詞：**取樣事件、鳥類、距離取樣、族群指標、公民科學

Received: October 05, 2017

Accepted: November 29, 2017

收件日期：2017年10月05日


接受日期：2017年11月29日

## Introduction

Information about the status of wildlife populations is vital for biodiversity conservation, especially as early warnings of emerging environmental crises and for evaluating the efficiency of implemented conservation strategies. Relevant datasets with a large spatial and temporal time scale are scarce in the Asia-Pacific region, including Taiwan. Wild bird populations are among the few taxa for which substantial data are available in Taiwan, mainly due to amateurs engaged in wildlife observation and bird counting events. Still, monitoring data collected by systematic sampling schemes are rare even for birds, possibly due to the financial and management challenges to run such programs. Although unstructured citizen-science

data are data rich and powerful in many circumstances, they may not be as sensitive as structured monitoring data to detect population changes (Kamp *et al.* 2016). Implementing a monitoring program that applies a systematic sampling scheme and also recruits the energy of citizen science hence seems promising and invaluable. For these reasons, the Taiwan Breeding Bird Survey project was initiated.

The Taiwan Breeding Bird Survey (BBS Taiwan) is the very first systematic regional-scale bird monitoring scheme in Taiwan and also one of the pioneers in the Asia-Pacific region. In 2009, a joint partnership by the Chinese Wild Bird Federation, the Institute of Ecology and Evolutionary Biology of National Taiwan University, and the Endemic Species Research Institute of the Council of Agriculture, Executive



Yuan, R.O.C. (Taiwan), was formed to initiate the project. Since then, the annual BBS Taiwan has recorded an average of 231 species (range 201-254) and 104,160 bird individuals (range 58,664 to 134,789) in 257 (range 140-355) sampling sites, with an average, 228 people (range 112-330) participating (Fig. 1). Running annually without an expected end of its project time, the BBS Taiwan is expected to continue to collect and update breeding bird population distribution and abundance data.

The primary aim of the project is to produce population indices of breeding birds in Taiwan to serve as biodiversity and environmental

indicators. To date, 102 breeding bird species population trends were reported (Ko *et al.* 2017). The usage of the dataset has broadened into a wide range of issues, including the Red-List of Birds of Taiwan (Lin *et al.* 2016), status and characteristics of introduced bird species in Taiwan (Su *et al.* 2014, 2015, 2016, 2017), and landscape planning (Huang 2015). Due to the standardized sampling protocol, comprehensive spatial coverage, and the increasing temporal coverage in the future, the usage of the dataset has even more potential in studying biogeography, community composition, and biodiversity indicators.




Fig. 1 Statistics of the BBS Taiwan from 2009 to 2015.

圖1. BBS Taiwan於2009至2015年的統計數據

## Project details

**Project title:** Bird sampling dataset of the Taiwan Breeding Bird Survey (BBS Taiwan)

**Funding:** Current funding from the Endemic

Species Research Institute, Council of Agriculture, Executive Yuan, R.O.C. (Taiwan). Initial funding was from the Forestry Bureau, Council of Agriculture, and the Biodiversity Research Center, National Taiwan University.



**Study area descriptions/descriptor:** As a nationwide monitoring project, the study extent is expected to cover both the Taiwan island and its main associated islands. This includes c. 36,190 square kilometers of land mass, located roughly between 21.5°N to 26.5°N and 118°E to 122°E. The climate is mainly subtropical with a mean temperature 18°C in winter and 28°C in summer, while average precipitation is 2,200 mm in the plains and over 3,000 mm in the mountains areas. Mountains compose 70% of the main

island, and the elevation ranges from 0 meters to 3,952 meters a.s.l..


## Taxonomic coverage

**General taxonomic coverage description:** A total of 313 bird species from 62 families have been recorded (Table 1), including all 27 endemic bird species of Taiwan, as well as most of the introduced bird species.


Table 1. List of bird orders and families recorded in the dataset

表1. 資料集內涵蓋的鳥類目別與科別

Order	Family	Family English Name
Anseriformes	Anatidae	Ducks, Geese, and Waterfowl
Galliformes	Phasianidae	Pheasants, Grouse, and Allies
Pelecaniformes	Pelecanidae	Pelicans
Pelecaniformes	Ardeidae	Hérons, Egrets, and Bitterns
Pelecaniformes	Threskiornithidae	Ibises and Spoonbills
Accipitriformes	Pandionidae	Osprey
Accipitriformes	Accipitridae	Hawks, Eagles, and Kites
Gruiformes	Rallidae	Rails, Gallinules, and Coots
Charadriiformes	Rostratulidae	Painted-Snipes
Charadriiformes	Jacanidae	Jacanas
Charadriiformes	Scolopacidae	Sandpipers and Allies
Charadriiformes	Turnicidae	Buttonquail
Charadriiformes	Laridae	Gulls, Terns, and Skimmers
Columbiformes	Columbidae	Pigeons and Doves
Cuculiformes	Cuculidae	Cuckoos
Strigiformes	Strigidae	Owls
Caprimulgiformes	Caprimulgidae	Nightjars and Allies
Caprimulgiformes	Apodidae	Swifts
Bucerotiformes	Bucerotidae*	Hornbills



Coraciiformes	Alcedinidae	Kingfishers
Coraciiformes	Coraciidae	Rollers
Piciformes	Megalaimidae	Asian Barbets
Piciformes	Picidae	Woodpeckers
Falconiformes	Falconidae	Falcons and Caracaras
Psittaciformes	Cacatuidae*	Cockatoos
Psittaciformes	Psittacidae*	New World and African Parrots
Passeriformes	Pittidae	Pittas
Passeriformes	Campephagidae	Cuckooshrikes
Passeriformes	Laniidae	Shrikes
Passeriformes	Vireonidae	Vireos, Shrike-Babblers, and Erpornis
Passeriformes	Oriolidae	Old World Orioles
Passeriformes	Dicruridae	Drongos
Passeriformes	Monarchidae	Monarch Flycatchers
Passeriformes	Corvidae	Crows, Jays, and Magpies
Passeriformes	Alaudidae	Larks
Passeriformes	Hirundinidae	Swallows
Passeriformes	Paridae	Tits, Chickadees, and Titmice
Passeriformes	Aegithalidae	Long-tailed Tits
Passeriformes	Troglodytidae	Wrens
Passeriformes	Cinclidae	Dippers
Passeriformes	Pycnonotidae	Bulbuls
Passeriformes	Cettiidae	Bush Warblers and Allies
Passeriformes	Regulidae	Kinglets
Passeriformes	Pnoepygidae	Cupwings
Passeriformes	Phylloscopidae	Leaf Warblers
Passeriformes	Acrocephalidae	Reed Warblers and Allies
Passeriformes	Locustellidae	Grassbirds and Allies
Passeriformes	Cisticolidae	Cisticolas and Allies
Passeriformes	Paradoxornithidae	Parrotbills, Wrentit, and Allies
Passeriformes	Zosteropidae	White-eyes, Yuhinas, and Allies
Passeriformes	Timaliidae	Tree-Babblers, Scimitar-Babblers, and Allies
Passeriformes	Pellorneidae	Ground Babblers and Allies
Passeriformes	Leiothrichidae	Laughingthrushes and Allies
Passeriformes	Muscicapidae	Old World Flycatchers



Passeriformes	Turdidae	Thrushes and Allies
Passeriformes	Sturnidae	Starlings
Passeriformes	Dicaeidae	Flowerpeckers
Passeriformes	Prunellidae	Accentors
Passeriformes	Motacillidae	Wagtails and Pipits
Passeriformes	Fringillidae	Finches, Euphonias, and Allies
Passeriformes	Passeridae	Old World Sparrows
Passeriformes	Estrildidae	Waxbills and Allies

\*introduced bird families

### Taxonomic ranks

Class: Ave

Common names: Bird

### Spatial coverage


**General spatial coverage:** Current records in the BBS Taiwan dataset are only from the main island of Taiwan and Lanyu Island off the east coast. The environmental gradient of the sampling locations spans an elevational range from 0–3,900m a.s.l. and across habitats from highly human-dominated urban landscapes to primary forests in remote mountain regions. Each sampling site was composed of 6–10 survey points with a 100 meter radius. Maximum distance between survey points at the same sites is 4 kilometers, and the minimum distance between survey points was 200 meters.

**Coordinates:** 21°48'36"N and 26°30'0"N Latitude; 118°22'48"E and 122°11'24"E Longitude

**Temporal coverage:** March 1, 2009 - July 31, 2015

### Methods

**Method step description:** The BBS Taiwan applied a stratified random sampling scheme for an initial set of 450 sampling sites aimed at covering 5% of the c.f. 36,000 km<sup>2</sup> area of Taiwan Island. Each sampling site is viewed as representative of a c.1 km x 1 km grid area, with survey points viewed as spatial repeats. Taiwan island is divided into 91 stratum, defined by 41 eco-regions (Su 1992) combined with 3 different elevational zones (0–1,000 m, 1,000–2,500 m, 2,500–4,000 m). We allocated the 450 sampling sites into each stratum by the area proportional to the area of each stratum. Additional sites, which we termed the “customized” set of samplings sites, were selected manually by surveyors at locations where no pre-selected sites existed. To date, sampling sites, including all types, number now approximately 762. All sampling sites were available to be adopted by one or several volunteer surveyors each year, whereby we encouraged the same surveyor to adopt the same sampling site as long as possible. Each sampling




sites was surveyed twice a year (once per trip), which, in combination with the 6–10 survey points per site, gives each site a minimum sampling effort of 12 repeated samples. The above sampling effort ensures the likelihood of detecting 80% of species present and keeping the coefficient of variance below 20 % in the breeding season within each site on Taiwan (Shiu and Lee 2003). Surveys are performed during the breeding season between March and June, with sites at higher elevations surveyed in a later sequence. Surveys are conducted within four hours after sunrise on each day. The point-count method (Bibby *et al.* 2000) with a 6–minute stop time was conducted on each survey point, ensuring that this level of sampling effort detected a minimum of 80% of the species at each site (Ding 1993, Shiu and Lee 2003). We record every detected bird individual, regardless

of migratory status or whether it was sighted or heard. For each bird record, the period of detection time (0–3 minutes, 3–6 minutes, and records beyond the 6–minute stop time are tagged as supplementary), horizontal distance to the observer (0–25m, 25–100m, >100m, flyover), and the formation of a flock with more than five individuals were recorded. For each survey point during each survey trip, the main-habitat, sub-habitat, presence of water-bodies, weather condition, and starting time were recorded. Habitats were classified by surveyors in the field according to categories listed in Table 2. Differences between the main and sub-habitat were judged by the percentage of area which each habitat type covers within a 100 meters radius, and only recording the two most prevalent types of habitats.


Table 2. BBS Taiwan habitat classification scheme

表2. BBS Taiwan棲地紀錄表

A-Forest	
A1	Broadleaved forest
A2	Conifer forest
A3	Mixed conifer-broadleaved forest (each >10%)
A4	Bamboo forest
A5	Mixed bamboo-broadleaved forest (each >10%)
A6	Windbreak forest (e.g. Casuarina stand)
A7	Other
B-Farmland	
B1	Rice field
B2	Other aquatic farmland (e.g. lotus or water chestnut field)
B3	Sugarcane field



B4	Dryland farm (e.g. vegetable, peanut or pineapple farm)
B5	Orchard or nursery (e.g. betel nut, tangerine, banana or wax apple orchard)
B6	Other
C-Grassland (natural and semi-natural)	
C1	Tall grassland (height >50cm, e.g. silvergrass grassland)
C2	Low grassland (height <50cm, e.g. the Lungpan grassland, pasture)
C3	High marsh (height >50cm, e.g. reed marsh)
C4	Low marsh (height <50cm, e.g. tuberous bulrush or sedge saltmarsh)
C5	Bamboo grassland
C6	Other
D-Inland (freshwater) wetland	
D1	Water storage area
D2	Lake (natural)
D3	River (water surface width >3m)
D4	Stream (water surface width <3m)
D5	Other
E-Aquaculture pond and saltpan	
E1	Flooded
E2	Dried
E3	Abandoned (vegetation overgrown to the center)
E4	Other
F-Coast	
F1	Tidal mudflat
F2	Tidal rock coast
F3	Rocky shoreline
F4	Sandy shoreline
F5	Marine
F6	Other
G-Urban and rural habitat	
G1	Urban building area
G2	Urban park and greenspace
G3	Rural setting
G4	Other
H-Other habitat type (with remarks)	



**Study extent description:** Sampling sites of the BBS Taiwan were selected to cover all major habitats on the main island of Taiwan, with an additional set on Lanyu Island that was surveyed during 2015. Other Taiwan's main associated islands is anticipated to be covered in the near future. Currently, the dataset contains

454 sampling sites, which includes a total of 3,901 survey points (Fig. 2). The major habitat types covered by these sampling sites include forests (48%), farmlands (24%), urban areas (12%), wetlands (7%), and grasslands (6%) (Fig. 3)




Fig 2. Distribution of sampling sites across the main island and Lanyu Island in Taiwan. Solid line on the right is the highest altitude in the survey area at each latitude.

圖2. 調查樣區於臺灣本島與蘭嶼的分布狀況；右圖實線為各緯度最高海拔。







Fig 3. Percentage of habitat type coverage of sampling sites.

圖3. 調查樣區的棲地分布比例

**Quality control description:** Efforts to maintain the quality of data were implemented before and after each survey. Before surveys, training workshops with field excursions were held to ensure that each volunteer surveyor fully understood the survey methods. After the data were submitted by the surveyors, the location of survey points, correctness of the time period, distance, and flock notes were reviewed the BBS Taiwan workgroup. Filters for rare or easily-misidentified species and unusually high numbers were also used to detect possibly erroneous

records. The highlighted records were then double-checked with surveyors, and doubtful records were deleted from the release version of the dataset. The taxonomic system follows the Checklist of Birds of Taiwan maintained by the Bird Record Committee of the Chinese Wild Bird Federation (Ding *et al.* 2017). All columns in the dataset are described in the metadata provided through the GBIF Integrated Publishing Toolkit (IPT) hosted by Taiwan Biodiversity Information Facility (TaiBIF). The dataset accompanying this data paper is the complete set



of the collected raw data.

## Datasets

### Dataset description

**Object name:** Darwin Core Archive The Taiwan Breeding Bird Survey Data

**Character encoding:** UTF-8

**Format name:** Darwin Core Archive format

**Format version:** 1.0

### Distribution:

[http://ipt.taibif.tw/archive.do?r=bbstaiwan\\_dataset](http://ipt.taibif.tw/archive.do?r=bbstaiwan_dataset)  
<https://www.gbif.org/dataset/f170f056-3f8a-4ef3-ac9f-4503cc854ce0>

**Publication date of data:** 2017-10-04

**Language:** English

**Licences of use:** This work is licensed under a Creative Commons Attribution (CC-BY) 4.0 License (<https://creativecommons.org/licenses/by/4.0/legalcode>).

**Metadata language:** English

**Date of metadata creation:** 2017-07-13

**Hierarchy level:** Dataset


## Acknowledgements

We thank all of the participants of BBS Taiwan, including the project organizers, regional organizers, workshop speakers, and survey volunteers. We thank Mai Guan-Shuo of the Systematics and Biodiversity Informatics Center with solving Darwin Core Archive difficulties, Liu Kuan-Ting for technical support in preparing the dataset, and An-Yu Chang for image production. The BBS Taiwan workgroup thanks

the Forestry Bureau, Council of Agriculture, and the Biodiversity Research Center, National Taiwan University, for funding the project in the initial stage. Bruno A. Walther edited this manuscript.

## References

- Bibby, C. J., N. D. Burgess, D. A. Hill, and S. H. Mustoe. 2000. Bird census techniques, 2nd edition. Academic, London, UK.
- Ding, T. S. 1993. Avian community ecology of mature forests on Mt. Yushan. Master Thesis. National Taiwan University, Taipei, Taiwan. (in Chinese, English abstract)
- Ding, T. S., C. S. Juan, R. S. Lin, C. Y. Pan, Y. J. Tsai, J. Wu, and Y. H. Yang. 2017. The 2017 CWBF checklist of the birds of Taiwan. Chinese Wild Bird Federation, Taipei, Taiwan.
- Huang, C. W. 2015. Developing a cell-based spatial optimization model for analyzing the trade-off between urban development and biological conservation. PhD Dissertation, National Taiwan University, Taipei, Taiwan.
- Kamp, J., S. Oppel, H. Heldbjerg, T. Nyegaard, and P. F. Donald. 2016. Unstructured citizen science data fail to detect long-term population declines of common birds in Denmark. *Diversity and Distribution* 22:1024–1035.
- Ko, C. J., M. W. Fan, Y. C. Lu, A. Y. Chang, L. F. Yen, R. Yang, Y. Y. Lo, R. S. Lin, S. P. Tsai, and P. F. Lee. 2017. 2015 Taiwan breeding bird survey annual report. Taiwan



- Endemic Species Research Institute, Nantou, Taiwan.
- Lin, R. S., Y. J. Lu., C. H. Yang, T. J. Tseng, C. J. Ko, and W. J. Chen. 2016. The red list of birds of Taiwan, 2016. Taiwan Endemic Species Research Institute, Nantou, Taiwan.
- Shiu, H. J., and P. F. Lee. 2003. Assessing avian point-count duration and sample size using species accumulation functions. *Zoological Studies* 42:357–367.
- Su, H. J. 1992. A geographical data organization system for the botanical inventory of Taiwan. Botany Institute, Academia Sinica Monograph Series 12:23-36.
- Su, S., P. Cassey, and T. Blackburn. 2014. Patterns of non-randomness in the composition and characteristics of the Taiwanese bird trade. *Biological Invasions* 16:2563-2575.
- Su, S., P. Cassey, and T. Blackburn. 2016. The wildlife pet trade as a driver of introduction and establishment in alien birds in Taiwan. *Biological Invasions* 18: 215–229
- Su, S., P. Cassey, M. Vall-Iloera, and T. Blackburn. 2015. Going cheap: Determinants of bird price in the Taiwanese pet market. *PloS one* 10:e0127482.
- Su, S., P. Cassey, E. E. Dyer, and T. Blackburn. 2017. Geographical range expansion of alien birds and environmental matching. *Ibis* 159: 193–203.

